Available online at www.sciencedirect.com
ScienceDirect
Nuclear Physics A 00 (2014) 1–4
Nuclear Physics A

Light flavor hadron spectra at low p_{T} and search for collective phenomena in high multiplicity pp, p–Pb and Pb–Pb collisions measured with the ALICE experiment

C. Andrei (for the ALICE Collaboration)

National Institute for Physics and Nuclear Engineering, Bucharest, Romania

arXiv:1408.0093v1 [hep-ex] 1 Aug 2014

Abstract

Comprehensive results on transverse momentum distributions and their ratios for identified light flavor hadrons (π, K, p) at low p_T and mid-rapidity as a function of charged particle multiplicity are reported for pp collisions at 7 TeV. Particle mass dependent hardening of the spectral shapes in Pb–Pb collisions at 2.76 TeV were attributed to hydrodynamical flow and quantitatively parameterized with Boltzmann-Gibbs Blast Wave fits. In this contribution, we investigate the existence of collective phenomena in small systems: pp, p–Pb and peripheral Pb–Pb where similar patterns are observed in multiplicity dependent studies.

Keywords: LHC, ALICE, light-flavor hadrons, multiplicity, transverse momentum spectra, collective phenomena

Identified charged particle transverse momentum spectra in p–pbar collisions were extensively studied as a function of incident energy below 900 GeV at CERN Sp$\bar{\mathrm{p}}$S [1] and up to 1800 GeV at Fermilab Tevatron [2]. Starting from 200 GeV, a deviation of the $\langle p_T \rangle$ of kaons as a function of $\sqrt{s}$ relative to the expectations based on the extrapolation of ISR data was reported by the UA5 Collaboration [3]. The E735 Collaboration saw evidence for a mass dependence of the $\langle p_T \rangle$ as a function of c.m. energy from 300 to 1800 GeV and of the $\langle p_T \rangle$ as a function of $\mathrm{d}N_{ch}/\mathrm{d}\eta$ at 1800 GeV [2]. The origin of these experimental observations is still under debate. QCD inspired models like PYTHIA [4] and EPOS [5] reproduce the experimental trends observed at Tevatron considering multiple partonic interactions and rescattering or a hydrodynamic type evolution with flux tube initial conditions, respectively. At about four times larger incident energies, the case of the present study, such processes become more important and contribute to a large energy transfer, well beyond the deconfinement energy density. Therefore, it is quite probable that at such energies a piece of matter of proton size explodes hydrodynamically once the energy transfer becomes significantly large [6].

Transverse momentum distributions and their relative ratios for identified positive charged hadrons π^+, K^+, p at mid-rapidity in pp collisions at $\sqrt{s}$ = 7 TeV, measured by the ALICE experiment [7] at the LHC, as a function of charged particle multiplicity are obtained. The p_{T} range goes from 0.2 GeV/c, 0.3 GeV/c and 0.5 GeV/c to 2.6 GeV/c, 1.4 GeV/c and 2.6 GeV/c for π^+, K^+ and p respectively, significantly larger than previously reported results [8]. 60 million inelastic pp collisions collected by the ALICE experiment during the 2010 run at LHC, using minimum bias (MB) and high multiplicity (HM) triggers [9], at $\sqrt{s}$ = 7 TeV, were analyzed. Events corresponding to the measured charged particle multiplicity within the pseudorapidity range $|\eta| < 0.8$ were selected. The multiplicity was estimated summing the global tracks and complementary Inner Tracking System (ITS) standalone tracks and Silicon Pixel Detector (SPD) tracklets. Events were required to have a primary vertex within ±10 cm from the center of the Time Projection Chamber (TPC) and the Time-Of-Flight array (TOF) subdetectors. A p_{T} dependent Distance of Closest



(a) (b)

Figure 1: a) Upper row - charged particle multiplicity dependence of the transverse momentum distributions for π^+, K^+ and p in pp collisions at 7 TeV; $z^{raw}=\left\langle N_{ch}^{raw}\right\rangle_{mult\ bin} / \left\langle N_{ch}^{raw}\right\rangle_{mult>0}$. Bottom row - ratio of transverse momentum distributions in a given multiplicity bin (z) relative to mult > 0; b) Upper row - p_T dependence of the particle ratios K^+/p, p/π^+ and p/K^+ in pp collisions at 7 TeV for two multiplicity bins. Bottom row - the ratio of the upper distributions relative to the one for mult > 0.

Approach (DCA) cut of $(0.018 + 0.035 p_T^{-1.01})$ cm in the transverse plane and 2 cm in the longitudinal direction were used in order to reduce the contamination from weak decays of strange particles, conversion or secondary hadronic interactions in the detector material. The present analysis has been done in $|y| < 0.5$. The particle identification was based on a Bayesian approach using the information from ITS, TPC and TOF. The detector response functions and the *a priori* probabilities were obtained from the experimental data. The contaminations from weak decays products and from particles produced by the interactions with the detector material were estimated based on a data driven method. All the correction factors were estimated using simulations based on the PYTHIA event generator [10] (tune Perugia0 [11]) and the GEANT3 [12] transport code. The corrections determined for the MB case were applied for all the multiplicity bins and their variations as a function of multiplicity, proved to be very small, were included in the systematic errors.

The fully corrected p_T spectra for π^+, K^+ and p were obtained by selecting events in eight multiplicity bins in the raw charged particle multiplicity distribution, 7-12, 13-19, 20-28, 29-39, 40-49, 50-59, 60-71 and 72-82 or, using the scaled multiplicity $z^{raw}=\left\langle N_{ch}^{raw}\right\rangle_{mult\ bin} / \left\langle N_{ch}^{raw}\right\rangle_{mult>0}$, [0.7-1.4), [1.4-2.1), [2.1-3.0), [3.0-4.2), [4.2-5.2), [5.2-6.3), [6.3-7.5) and [7.5-8.6]. The results are presented in Fig. 1a - upper row. In the bottom row of Fig. 1a the ratios of p_T distributions at different multiplicities relative to the mult > 0 case are represented. One could observe a systematic change in the shape of the ratios, i.e. a depletion in the low p_T region which shows a tendency to level off at larger p_T values. The amount of depletion clearly depends on the mass of the species and on the multiplicity, i.e. it is enhanced going from pions to protons and with increasing multiplicity for a given mass. The K^+/π^+, p/π^+ and p/K^+ ratios as a function of p_T are plotted in Fig. 1b - upper row for the second and the sixth multiplicity bins. The general trends observed in Fig. 1a can be seen in a more quantitative way in these representations, especially in the bottom row of Fig. 1b, where the ratios of relative yields for the two multiplicities to the ones corresponding to $N_{ch}^{raw} > 0$ are represented. The observed depletion of p/π^+ and p/K^+ at low p_T, increasing with the multiplicity and decreasing towards 1.4-2.0 GeV/c looks similar with the trends observed in A–A collisions, the heavier particles being pushed towards larger transverse momenta. Such behavior, observed in Pb–Pb collisions at 2.76 TeV [13], was attributed to



(a) (b) (c)

Figure 2: a) p_{T} dependence of p/π^+ ratio for the second and highest multiplicity bins in pp collisions; b) p_{T} dependence of $(\mathrm{p}+\overline{\mathrm{p}})/(\pi^+ + \pi^-)$ ratio in p–Pb for 60-80% and 0-5% multiplicity classes and Pb–Pb for 80-90% and 0-5% centrality [14]; c) The ratios of the ratios presented in a) and b).

the existence of collective transverse flow. The evolution of the p_{T} spectra shape with charged particle multiplicity in pp collisions at 7 TeV and p–Pb at 5.02 TeV [14] and with centrality in Pb–Pb at 2.76 TeV [13] is rather similar.

The p_{T} dependence of p/π^+ for the second and highest multiplicity bins and of $(\mathrm{p}+\overline{\mathrm{p}})/(\pi^+ + \pi^-)$ ratio for p–Pb in 60-80% and 0-5% multiplicity classes and for Pb–Pb at 80-90% and 0-5% centrality [14] are presented in Fig. 2a and Fig. 2b, respectively. The push of protons towards larger p_{T} values relative to pions with increasing centrality or multiplicity is present for all three systems. Quantitatively, this can be followed in Fig. 2c where the ratios of the ratios shown in Fig. 2a and Fig. 2b are presented. The ratio for pp follows closely the p–Pb trend as a function of p_{T}.

Based on these similarities one could investigate to what extent the quality and parameters of simultaneous fits of transverse momentum distributions for different multiplicities or centralities using expressions inspired by hydrodynamic models are also similar. The quality of fits for π^+, K^+ and p, based on the Boltzman-Gibbs Blast Wave (BGBW) expression [15]:

$$E\frac{d^3N}{dp^3} \sim f(p_T) = \int_0^R m_T K_1(m_T cosh\rho/T_{kin}) I_0(p_T sinh\rho/T_{kin}) r dr \quad (1)$$

where $m_T = \sqrt{m^2 + p_T^2}$; $\beta_r(r) = \beta_s(\frac{r}{R})^n$; $\rho = tanh^{-1}\beta_r$, is similar within the error bars for all three systems. The p_{T} range on which the p_{T} spectra follows the hydro shape increases going towards higher multiplicity in pp and p–Pb or higher centrality in Pb–Pb.

In Fig. 3a the results of the fits for pp collisions at 7 TeV in terms of kinetic freeze-out temperature (T_{kin}) versus transverse expansion velocity ($\langle\beta_T\rangle$) are presented for different multiplicity bins and compared with the results obtained for p–Pb and Pb–Pb as a function of multiplicity classes and centrality, respectively [14]. While the T_{kin}-$\langle\beta_T\rangle$ correlation as a function of multiplicity in pp overlaps with the one corresponding to p–Pb as a function of multiplicity classes, for Pb–Pb the decrease of T_{kin} and increase of $\langle\beta_T\rangle$ with centrality is more enhanced. The observed trends are not reproduced by PYTHIA in absolute values although a somewhat similar trend seems to be present if Color Reconnection is included.

Further information accessed from these fits is the expansion profile, given by the parameter n. The n-$\langle\beta_T\rangle$ correlation as a function of multiplicity or centrality for the three systems is presented in Fig. 3b. Towards high charged particle multiplicity, the transverse expansion velocity approaches a linear dependence as a function of position in the fireball for pp, the trend overlapping with the one corresponding to p–Pb and Pb–Pb.

In conclusion, transverse momentum spectra of positive identified charged hadrons as a function of charged particle multiplicity in pp collisions at $\sqrt{s}$ = 7 TeV, up to multiplicity values never measured before, are reported. A clear depletion of p/π^+ and p/K^+ at low p_{T} , increasing with the multiplicity and decreasing towards larger p_{T}, similar with the trends observed in p–Pb and Pb–Pb collisions, is seen. The kinetic freeze-out temperature (T_{kin}), expansion


T_{kin} (GeV)
ALICE
PRELIMINARY
ALICE, p-Pb, $\sqrt{s_{NN}}$ = 5.02 TeV
V0A Multiplicity Classes (Pb-side)
ALICE, Pb-Pb, $\sqrt{s_{NN}}$ = 2.76 TeV
ALICE, pp, $\sqrt{s}$ = 7 TeV
PYTHIA8, $\sqrt{s}$ = 7 TeV (with Color Reconnection)
PYTHIA8, $\sqrt{s}$ = 7 TeV (without Color Reconnection)
$\langle\beta_T\rangle$
n
ALICE Preliminary
pp, $\sqrt{s}$ = 7 TeV
p-Pb, $\sqrt{s_{NN}}$ = 5.02 TeV
Pb-Pb, $\sqrt{s_{NN}}$ = 2.76 TeV
Uncertainties: syst.(from global fit of data)
Fit-range related syst. uncert.- not included
<β_T>

(a) (b)

Figure 3: a) Freeze-out T_{kin} versus transverse expansion velocity $\langle\beta_T\rangle$ for different multiplicity bins for pp collisions at 7 TeV (black triangles) comparison with the results obtained for p–Pb (dark blue dots) for different multiplicity classes and Pb–Pb (red circles) as a function of centrality [14]; b) Expansion profile n versus transverse expansion velocity $\langle\beta_T\rangle$ for different multiplicity bins for pp collisions at 7 TeV (green dots) compared with the results obtained for p–Pb (dark blue triangles) dots for different multiplicity classes and Pb–Pb (red squares) as a function of centrality [14]. The $\langle\beta_T\rangle$ values increase with multiplicity (pp and p–Pb) or centrality (Pb–Pb).

velocity ($\langle\beta_T\rangle$) and its profile extracted from the simultaneous fits of the π^+, K^+ and p spectra with BGBW, show a multiplicity dependence trend similar with the ones obtained for p–Pb and Pb–Pb collisions as a function of multiplicity classes or centrality, respectively. However, a conclusion about similar mechanisms for the three systems has to be taken with caution. Detailed investigations based on theoretical approaches such as hydrodynamic models, parton based Gribov-Regge theory, Color Glass Condensate, Color Reconnection, will give insight to the underlying physics of this similar behavior observed at LHC energies.

References

[1] R.E.Ansorge et al., Phys.Lett.B, 167, (1986), p. 476.
[2] T.Alexopoulos et al., Phys.Rev.D, 48, (1993), p. 984.
[3] G.J.Alner et al., Nucl.Phys.B, 258, (1985), p. 505.
[4] T.Sjöstrand and P.Z.Skands, arXiv:hep-ph/0402078; R.Corke, arXiv:hep-ph/o901.2852.
[5] K.Werner et al., Phys.Rev.C, 74, (2006), p. 044902.
[6] S.Z.Belenkji, L.D. Landau, Del Nuov.Cim., Suppl. Vol.III, Serie X, (1956), p. 15; G.A.Malekhin, Zh.Eksp.Teor.Fiz., 35, (1958), p. 1185; L.Van Hove, Phys.Lett.B, 118, (1982), p. 138; B.Andersson et al., Physica Scripta, 20, (1979), p. 10.
[7] K.Aamodt et al., Eur. Phys. J. C, 68, (2010), p. 345.
[8] CMS Collaboration, Eur. Phys. J. C, 72, (2012), p. 2164.
[9] ALICE Collaboration, arXiv:nucl-ex/1402.4476.
[10] T. Sjöstrand, S.Mrenna, P.Z.Skands, arXiv:hep-ph/0603175.
[11] M.G.Albrow et al., arXiv:hep-ph/0610012.
[12] R.Brun, F.Carminati, S.Giani, CERN-W5013.
[13] B. Abelev et al., Phys.Rev.C, 88, (2013), p. 044910.
[14] B. Abelev et al., Phys.Lett.B, 728, (2014), p. 25.
[15] E.Schnedermann et al., Phys.Rev.C, 48, (1993), p. 2462.